

***Phaeomonilia aquatica* sp. nov., an aquatic hyphomycete from China**

JUN-EN HUANG¹, HAI-YAN SONG², XI-GEN HUANG³,
JIAN MA^{2*} & DIAN-MING HU^{1*}

¹ Jiangxi Key Laboratory for Conservation and Utilization of Fungal Resources,

² College of Agronomy, Jiangxi Agricultural University & ³ College of Science,
Jiangxi Agricultural University, Nanchang, 330045, China

* CORRESPONDENCE TO: Jian Ma, Dian-Ming Hu, email: hudianming1@163.com

ABSTRACT—A new aquatic hyphomycete, *Phaeomonilia aquatica*, is described and illustrated from specimens collected on submerged wood in Jiangxi Province, China. *Phaeomonilia aquatica* is morphologically characterized by conidiophores with percurrent proliferation and aseptate smooth conidia that are spherical, subtriangular, subglobose, ellipsoidal to subfusiform or obovate and formed in branched acropetal chains. A key to *Phaeomonilia* species is provided.

KEY WORDS — anamorphic fungi, freshwater fungi, lignicolous fungi, taxonomy

Introduction

Hyphomycetous fungi are highly diverse in freshwater habitats (Hu et al. 2013). As essential components in aquatic ecosystems, they play an irreplaceable role in the natural cycle of organic materials and energy. Their global distribution is poorly studied (Shearer et al. 2007), but our group has investigated the distribution of aquatic hyphomycetes for about ten years (Hu et al. 2007, 2010a,b, 2012a,b,c, 2013; Huang et al. 2016). The ongoing investigation in Jiangxi Province, China, has led to the collection of a new species of *Phaeomonilia* R.F. Castañeda et al.

Materials & methods

Unidentified wood samples submerged in a freshwater stream were collected in Jiangxi Province, China, and incubated in moist chambers at room temperature (ca. 25

°C). Samples were examined for fungal fruiting bodies using a dissecting microscope. Observations and photographs were prepared from materials mounted in water and examined with a Nikon Ni compound microscope (Hu et al. 2012a). The specimens examined were deposited in the Herbarium of Fungi, Jiangxi Agricultural University, Nanchang, China (HFJAU).

Taxonomy

Phaeomonilia aquatica J.E. Huang, H.Y. Song, Jian Ma & D.M. Hu, **sp. nov.** PL1

MYCOBANK MB 821142

Differs from *Phaeomonilia guangxiensis* by its conidiophores with percurrent proliferations through the apex and by its narrower conidia, and from *P. nanningensis* by its conidiophores with percurrent proliferations through the apex and by its shorter hyaline to yellowish conidia.

TYPE—China, Jiangxi Province, Jiujiang, Lushan Mountain, in a stream on submerged wood, 13 Oct. 2016, J.E. Huang (**Holotype**, HFJAU 0148).

ETYMOLOGY—*aquatica*, referring to the aquatic habitat of the fungus.

COLONIES on submerged wood effuse, brown to dark brown, consisting of scattered single conidiophores over the substrate surface. MYCELIUM immersed or superficial, composed of brown, cylindrical, septate, infrequently branched hyphae. CONIDIOPHORES macronematous, mononematous, sometimes more than 900 µm long and 8.5–18 µm diam., single, erect, straight or flexuous, branched or unbranched, smooth, occasionally rough, 15–35-septate, thin-walled, brown to dark brown toward the base, fading slightly toward the apex, with percurrent proliferations through the apex, giving rise to new conidiogenous loci at the apex. CONIDIOGENOUS CELLS holothallic, 12–29 × 8.5–13 µm, terminal, subapical or lateral, obovoid, cupulate or obtrapezoid, smooth, pale brown, with several pale brown to hyaline trapezoid denticles, 3.0–5.9 × 0.9–2.3 µm. CONIDIA thallic-arthric, 1-celled, 8–14 × 7.5–12 µm, spherical, subtriangular, subglobose, ellipsoidal to subfusiform or obovate, hyaline to yellowish, in branched acropetal chains of about seven conidia, thin-walled, with some small wart on the surface. Conidial secession schizolytic.

ADDITIONAL SPECIMEN EXAMINED—CHINA, JIANGXI PROVINCE, Jiujiang, Lushan Mountain, in a stream on submerged wood, 13 Oct. 2016, J.-E. Huang (HFJAU 0323).

PLATE 1. *Phaeomonilia aquatica* (holotype, HFJAU 0148). a. Conidiophore and conidia on submerged wood; b–d. Conidiophores and conidia. Note the proliferation (arrowed in d); e–f. Conidial chains; g. Conidiogenous cell with conidia; h–k. Conidiogenous cells; l–s. Conidia. Scale bars: a, b = 100 µm; c–f = 50 µm; g = 20 µm; h–s = 5 µm.



Discussion

Castañeda-Ruíz et al. (2007) erected *Phaeomonilia* for two species, *P. pleiomorpha* R.F. Castañeda et al. (type species) and *P. corticola* (R.F. Castañeda) R.F. Castañeda et al. [≡ *Monilia corticola* R.F. Castañeda]. They compared *Phaeomonilia* morphologically with similar genera, including *Botryomonilia* Goos & Piroz., *Staheliella* Emden, and *Oidiodendron* Robak. Subsequently, Ma et al. (2015) and Yang et al. (2016) described *P. guangxiensis* Y.R. Ma & X.G. Zhang and *P. nanningensis* C.L. Yang & X.G. Zhang from China. *Phaeomonilia pleiomorpha* was collected from a palm petiole submerged in a stream, but the other three species were collected from decaying petioles, twigs, or stems in terrestrial habitats. *Phaeomonilia aquatica* is the second species reported from freshwater habitats.

Phaeomonilia aquatica can be easily distinguished from the other species of *Phaeomonilia* by its percurrently proliferating conidiophores. In addition, *P. pleiomorpha* differs from *P. aquatica* by its smaller, globose, broadly Y-shaped conidia ($4.0\text{--}5.8 \times 3.9\text{--}4.4\text{ }\mu\text{m}$) and its *Stylaspergillus*-like synanamorph (Castañeda-Ruíz et al. 2007); *P. guangxiensis* differs by its slightly shorter and wider, colorless, ellipsoid to globose, broadly Y-shaped conidia ($7\text{--}11 \times 11\text{--}14.5\text{ }\mu\text{m}$; Ma et al. 2015); *P. nanningensis* differs by its longer and slightly wider, brown to pale brown conidia ($13.5\text{--}25 \times 9\text{--}13\text{ }\mu\text{m}$; Yang et al. 2016); and *P. corticola* differs by its narrower, doliiform or ellipsoid conidia ($5.5\text{--}16 \times 4\text{--}5\text{ }\mu\text{m}$; Castañeda-Ruíz et al. 2007). A key to *Phaeomonilia* species is presented below.

Key to species of *Phaeomonilia*

- 1. *Stylaspergillus*-like synanamorph present *P. pleiomorpha*
- 1. No *Stylaspergillus*-like synanamorph 2
- 2. Conidia doliiform or ellipsoid, $5.5\text{--}16 \times 4\text{--}5\text{ }\mu\text{m}$ *P. corticola*
- 2. Conidia not doliiform, $>7.5\text{ }\mu\text{m}$ diam. 3
- 3. Conidiophores with percurrent proliferation; conidia spherical, subtriangular, subglobose, ellipsoidal to subfusiform, or obovate. *P. aquatica*
- 3. Conidiophores without percurrent proliferation; conidia ellipsoidal, globose, or Y-shaped4
- 4. Conidia hyaline, $7\text{--}11 \times 11\text{--}14.5\text{ }\mu\text{m}$ *P. guangxiensis*
- 4. Conidia brown to pale brown, $13.5\text{--}25 \times 9\text{--}13\text{ }\mu\text{m}$ *P. nanningensis*

Acknowledgments

The authors express their sincere gratitude to Dr. Eric H.C. McKenzie and Dr. Huang Zhang for their critical review of the manuscript, to Dr. Shaun Pennycook for

nomenclatural review, and to Dr. Lorelei L. Norvell for editorial review. This project was supported by the National Natural Science Foundation of China (Nos. 31500021, 31460009) and The Key Research and Development Program of Jiangxi Province (No. 20161BBF60078).

Literature cited

- Castañeda-Ruiz RF, Heredia-Abarca G, Arias RM, Saikawa M, Minter DW, Stadler M. 2007. Anamorphic fungi from submerged plant material: *Phaeomonilia pleiomorpha*, *P. corticola* and *Cacumisporium pleuroconidiophorum*. *Mycotaxon* 100: 327–336.
- Hu DM, Zhu H, Cai L, Hyde KD, Zhang KQ. 2007. *Sirothecium triseriale*, a new chirosporous anamorphic species from China. *Cryptogamie, Mycologie* 28: 311–314.
- Hu DM, Cai L, Chen H, Bahkali AH, Hyde KD. 2010a. Four new freshwater fungi associated with submerged wood from Southwest Asia. *Sydowia* 62: 191–203.
- Hu DM, Cai L, Chen H, Bahkali AH, Hyde KD. 2010b. Fungal diversity on submerged wood in a tropical stream and an artificial lake. *Biodiversity and Conservation* 19: 3799–3808. <https://doi.org/10.1007/s10531-010-9927-5>
- Hu DM, Cai L, Bahkali AH, Hyde KD. 2012a. Two new freshwater species of *Annulatascaceae* from China. *Mycotaxon* 120: 81–88. <https://doi.org/10.5248/120.81>
- Hu DM, Cai L, Hyde KD. 2012b. Three new ascomycetes from freshwater in China. *Mycologia* 104: 1478–1489. <https://doi.org/10.3852/11-430>
- Hu DM, Chen H, Cai L, Bahkali AH, Hyde KD. 2012c. *Aquapeziza*: a new genus from freshwater, and its morphological and phylogenetic relationships to *Pezizaceae*. *Mycologia* 104: 540–546. <https://doi.org/10.3852/11-123>
- Hu DM, Liu F, Cai L. 2013. Biodiversity of aquatic fungi in China. *Mycology* 4: 125–168. <https://doi.org/10.1080/21501203.2013.835752>
- Huang JE, Song HY, Ma J, Guan GX, Hu DM. 2016. *Xylohyphopsis aquatica* sp. nov., a new aquatic hyphomycete from China. *Mycotaxon* 131: 391–394. <https://doi.org/10.5248/131.391>
- Ma YR, Xia JW, Zhang XG, Castañeda-Ruiz RF. 2015. New species of *Phaeomonilia* and *Mirandina* from dead branches in China. *Mycotaxon* 130: 775–781. <https://doi.org/10.5248/130.775>
- Shearer CA, Descals E, Kohlmeyer B, Kohlmeyer J, Marvanová L, Padgett D, Porter D, Raja HA, Schmit JP, Thorton HA, Voglmayr H. 2007. Fungal biodiversity in aquatic habitats. *Biodiversity and Conservation* 16: 49–67. <https://doi.org/10.1007/s10531-006-9120-z>
- Yang CL, Wang JY, Xia JW, Ma YR, Gao JM, Zhang XG. 2016. *Phaeomonilia nanningensis* sp. nov. and a new *Craspedodidymum* record from southern China. *Mycotaxon* 131: 547–551. <https://doi.org/10.5248/131.547>